

NCC

*NNP-BC
Neonatal Nurse Practitioner*

For More Information – Visit link below:

<https://www.examsempire.com/>

Product Version

- 1. Up to Date products, reliable and verified.**
- 2. Questions and Answers in PDF Format.**



<https://examsempire.com/>

Latest Version: 6.0

Question: 1

most commonly identified trisomy in abortuses (30% of all autosomal aneuploidy in abortuses).

- A. Trisomy 16
- B. Trisomy 2
- C. Trisomy 21
- D. Trisomy 8

Answer: A

Question: 2

Trough <10
Peak 20-40

- A. Vancomycin levels
- B. Antibiotic length of therapy
- C. Calorie goals
- D. Weight classifications

Answer: A

Question: 3

increased pancreatic cell stimulation, or decreased glycolysis

- A. S/S urea cycle defects
- B. The mechanism by which maternal sympathomimetic drugs cause hypoglycemia is
- C. Anion Gap
- D. Beckwith-Wiedemann syndrome

Answer: B

Question: 4

Deposits of Ca++ in renal parenchyma, r/t PTM, loop diuretics, increased Ca++/VD supps, asymptomatic

- A. Catecholamines
- B. Staphylococci
- C. Ampicillin
- D. Nephrocalcinosis

Answer: D

Question: 5

$(\text{rate/hr})(\% \text{dex})(.167)/\text{wt in kgs}$
Advance to 8-12 mcg/kg/min for nutrition as needed

- A. ANGUS
- B. BRAHMAN
- C. GUZERAT
- D. GIR

Answer: D

Question: 6

Toxoplasmosis
Other
Rubella
Cytomegalovirus
Herpes

- A. Cytomegalovirus
- B. Gonorrhea
- C. Cystic fibrosis
- D. TORCH infections

Answer: D

Question: 7

Constant flow in pulmonary artery on doppler

- A. TETRALOGY OF FALLOT
- B. TRICUSPID ATRESIA
- C. PDA

D. COARCTATION OF THE AORTA

Answer: C

Question: 8

ACE, Alcohol, caffeine, carbamazepine, DES, phenytoin, Retin A, lithium, phenobarbital, salicylate, tetracycline, valproic acid, warfarin

- A. Tocolytics
- B. Carcinogens
- C. Mutagens
- D. Teratogens

Answer: D

Question: 9

Na 135-145
Cl 97-110
CO₂ 20-28
K 3.5-5.5
Mg 0.3-1
Ca 8-11 (iCa 50%)

- A. Aldosterone
- B. Thrombosis
- C. Carbohydrates
- D. Electrolytes

Answer: D

Question: 10

microcephaly, periventricular Ca, petechiae with thrombocytopenia, sensorineural hearing loss, blueberry muffin rash
urine or saliva CMV culture: neg exculde
serum CMV IgM antibody suggests
Herpes virus family (DNA)
Most common congenital infx
Get urine cultures within 1st week of life
Ganciclovir, platelets

80-90% asymptomatic

- A. SYPHILIS
- B. HEP B
- C. CHLAMYDIA
- D. CMV

Answer: D

Question: 11

Aminoglycoside prototype: bactericidal inhibitor of protein synthesis (30S); active against many aerobic gram-negative bacteria

a. Narrow therapeutic window; dose reduction required in renal impairment. Tox: renal dysfunction, ototoxicity; once-daily dosing is effective (post-antibiotic effect) and less toxic

- A. Clindamycin
- B. Vancomycin
- C. Gentamicin
- D. Ampicillin

Answer: C

Question: 12

Threshold maintained in proximal tubule, once reached, excreted in urine

- A. Bicarb
- B. GIR
- C. Dopamine
- D. Lasix

Answer: A

Thank You for Trying Our Product

Special 16 USD Discount Coupon: **NSZUBG3X**

Email: support@examsempire.com

**Check our Customer Testimonials and ratings
available on every product page.**

Visit our website.

<https://examsempire.com/>