

# Nursing

NCSBN-NCLEX-PN

ational Council of State Boards of Nursing: National Council Licensure Examination-Practical Nurse

For More Information – Visit link below:

<https://www.examsempire.com/>

**Product Version**

1. Up to Date products, reliable and verified.
2. Questions and Answers in PDF Format.



<https://examsempire.com/>

---

# Latest Version: 6.0

## Question: 1

A middle-aged woman tells the nurse that she has been experiencing irregular menses for the past six months. The nurse should assess the woman for other symptoms of:

- A. climacteric.
- B. menopause.
- C. perimenopause.
- D. postmenopause.

**Answer: C**

Explanation:

Perimenopause refers to a period of time in which hormonal changes occur gradually, ovarian function diminishes, and menses become irregular. Perimenopause lasts approximately five years. Climacteric is a term applied to the period of life in which physiologic changes occur and result in cessation of a woman's reproductive ability and lessened sexual activity in males. The term applies to both genders. Climacteric and menopause are interchangeable terms when used for females. Menopause is the period when permanent cessation of menses has occurred. Postmenopause refers to the period after the changes accompanying menopause are complete.

Health Promotion and Maintenance

## Question: 2

When obtaining a health history on a menopausal woman, which information should a nurse recognize as a contraindication for hormone replacement therapy?

- A. family history of stroke
- B. ovaries removed before age 45
- C. frequent hot flashes and/or night sweats
- D. unexplained vaginal bleeding

**Answer: D**

Explanation:

Unexplained vaginal bleeding is a contraindication for hormone replacement therapy. Family history of stroke is not a contraindication for hormone replacement therapy. If the woman herself had a history of stroke or other blood-clotting events, hormone therapy could be contraindicated. Frequent hot flashes and/or night sweats can be relieved by hormone replacement therapy.

Health Promotion and Maintenance

### Question: 3

Which of the following statements, if made by the parents of a newborn, does not indicate a need for further teaching about cord care?

- A. "I should put alcohol on my baby's cord 3–4 times a day."
- B. "I should put the baby's diaper on so that it covers the cord."
- C. "I should call the physician if the cord becomes dark."
- D. "I should wash my hands before and after I take care of the cord."

**Answer: D**

Explanation:

Parents should be taught to wash their hands before and after providing cord care. This prevents transferring pathogens to and from the cord. Folding the diaper below the cord exposes the cord to air and allows for drying. It also prevents wet or soiled diapers from coming into contact with the cord. Current recommendations include cleaning the area around the cord 3–4 times a day with a cotton swab but do not include putting alcohol or other antimicrobials on the cord. It is normal for the cord to turn dark as it dries. Health Promotion and Maintenance

### Question: 4

The nurse is teaching parents of a newborn about feeding their infant. Which of the following instructions should the nurse include?

- A. Use the defrost setting on microwave oven to warm bottles.
- B. When refrigerating formula, don't feed the baby partially used bottles after 24 hours.
- C. When using formula concentrate, mix two parts water and one part concentrate.
- D. If a portion of one bottle is left for the next feeding, go ahead and add new formula to fill it.

**Answer: A**

Explanation:

Parents must be careful when warming bottles in a microwave oven because the milk can become superheated. When a microwave oven is used, the defrost setting should be chosen, and the temperature of the formula should be checked before giving it to the baby. Refrigerated, partially used bottles should be discarded after 4 hours because the baby might have introduced some pathogens into the formula. Returning the bottle to the refrigerator does not destroy pathogens. Formula concentrate and water are usually mixed in a 1:1 ratio of one part concentrate and one part water. Infants should be offered fresh formula at each feeding. Partially used bottles should not have fresh formula added to them. Pathogens can grow in partially used bottles of formula and be transferred to the new formula. Health Promotion and Maintenance

### Question: 5

---

The nurse is assessing the dental status of an 18-month-old child. How many teeth should the nurse expect to examine?

- A. 6
- B. 8
- C. 12
- D. 16

**Answer: C**

Explanation:

In general, children begin dentition around 6 months of age. During the first 2 years of life, a quick guide to the number of teeth a child should have is as follows: Subtract the number 6 from the number of months in the age of the child. In this example, the child is 18 months old, so the formula is  $18 - 6 = 12$ . An 18-month-old child should have approximately 12 teeth.

Health Promotion and Maintenance

### Question: 6

Which of the following physical findings indicates that an 11–12-month-old child is at risk for developmental dysplasia of the hip?

- A. refusal to walk
- B. not pulling to a standing position
- C. negative Trendelenburg sign
- D. negative Ortolani sign

**Answer: B**

Explanation:

The nurse might be concerned about developmental dysplasia of the hip if an 11–12-month-old child doesn't pull to a standing position. An infant who does not walk by 15 months of age should be evaluated. Children should start walking between 11–15 months of age. Trendelenburg sign is related to weakness of the gluteus medius muscle, not hip dysplasia. Ortolani sign is used to identify congenital subluxation or dislocation of the hip in infants.

Health Promotion and Maintenance

### Question: 7

When administering intravenous electrolyte solution, the nurse should take which of the following precautions?

- A. Infuse hypertonic solutions rapidly.
- B. Mix no more than 80 mEq of potassium per liter of fluid.
- C. Prevent infiltration of calcium, which causes tissue necrosis and sloughing.

D. As appropriate, reevaluate the client's digitalis dosage. He might need an increased dosage because IV calcium diminishes digitalis's action.

**Answer: C**

Explanation:

Preventing tissue infiltration is important to avoid tissue necrosis. Choice 1 is incorrect because hypertonic solutions should be infused cautiously and checked with the RN if there is a concern. Choice 2 is incorrect because potassium, mixed in the pharmacy per physician order, is mixed at a concentration no higher than 60 mEq/L.

Physiological Adaptation

### Question: 8

Teaching about the need to avoid foods high in potassium is most important for which client?

- A. a client receiving diuretic therapy
- B. a client with an ileostomy
- C. a client with metabolic alkalosis
- D. a client with renal disease

**Answer: D**

Explanation:

Clients with renal disease are predisposed to hyperkalemia and should avoid foods high in potassium. Choices 1, 2, and 3 are incorrect because clients receiving diuretics with ileostomy or with metabolic alkalosis are at risk for hypokalemia and should be encouraged to eat foods high in potassium.

Physiological Adaptation

### Question: 9

What do the following ABG values indicate: pH 7.38, PO<sub>2</sub> 78 mmHg, PCO<sub>2</sub> 36mmHg, and HCO<sub>3</sub> 24 mEq/L?

- A. metabolic alkalosis
- B. homeostasis
- C. respiratory acidosis
- D. respiratory alkalosis

**Answer: B**

Explanation:

These ABG values are within normal limits. Choices 1, 3, and 4 are incorrect because the ABG values indicate none of these acid-base disturbances.

Physiological Adaptation

### Question: 10

The major electrolytes in the extracellular fluid are:

- A. potassium and chloride.
- B. potassium and phosphate.
- C. sodium and chloride.
- D. sodium and phosphate.

**Answer: C**

Explanation:

Sodium and chloride are the major electrolytes in the extracellular fluid.

Physiological Adaptation

### Question: 11

A client with Kawasaki disease has bilateral congestion of the conjunctivae, dry cracked lips, a strawberry tongue, and edema of the hands and feet followed by desquamation of fingers and toes. Which of the following nursing measures is most appropriate to meet the expected outcome of positive body image?

- A. administering immune globulin intravenously
- B. assessing the extremities for edema, redness and desquamation every 8 hours
- C. explaining progression of the disease to the client and his or her family
- D. assessing heart sounds and rhythm

**Answer: C**

Explanation:

Teaching the client and family about progression of the disease includes explaining when symptoms can be expected to improve and resolve. Knowledge of the course of the disease can help them understand that no permanent disruption in physical appearance will occur that could negatively affect body image. Clients with Kawasaki disease might receive immune globulin intravenously to reduce the incidence of coronary artery lesions and aneurysms. Cardiac effects could be linked to body image, but Choice 3 is the most direct link to body image. The nurse assesses symptoms to assist in evaluation of treatment and progression of the disease.

Health Promotion and Maintenance

### Question: 12

Which of the following is most likely to impact the body image of an infant newly diagnosed with Hemophilia?

- A. immobility
- B. altered growth and development
- C. hemarthrosis
- D. altered family processes

**Answer: D**

Explanation:

Altered Family Processes is a potential nursing diagnosis for the family and client with a new diagnosis of Hemophilia. Infants are aware of how their caregivers respond to their needs. Stresses can have an immediate impact on the infant's development of trust and how others relate to them because of their diagnosis. The longterm effects of hemophilia can include problems related to immobility. Altered growth and development could not have developed in a newly diagnosed client. Hemarthrosis is acute bleeding into a joint space that is characteristic of hemophilia. It does not have an immediate effect on the body image of a newly diagnosed hemophiliac.

Health Promotion and Maintenance

### Question: 13

While undergoing fetal heart monitoring, a pregnant Native-American woman requests that a medicine woman be present in the examination room. Which of the following is an appropriate response by the nurse?

- A. "I will assist you in arranging to have a medicine woman present."
- B. "We do not allow medicine women in exam rooms."
- C. "That does not make any difference in the outcome."
- D. "It is old-fashioned to believe in that."

**Answer: A**

Explanation:

This statement reflects cultural awareness and acceptance that receiving support from a medicine woman is important to the client. The other statements are culturally insensitive and unprofessional.

Reduction of Risk Potential

### Question: 14

All of the following should be performed when fetal heart monitoring indicates fetal distress except:

- A. increase maternal fluids.
- B. administer oxygen.
- C. decrease maternal fluids.
- D. turn the mother.

**Answer: C**

---

Explanation:

Decreasing maternal fluids is the only intervention that should not be performed when fetal distress is indicated.

Reduction of Risk Potential

### Question: 15

Which fetal heart monitor pattern can indicate cord compression?

- A. variable decelerations
- B. early decelerations
- C. bradycardia
- D. tachycardia

**Answer: A**

Explanation:

Variable decelerations can be related to cord compression. The other patterns are not.

Reduction of Risk Potential



## Thank You for Trying Our Product

Special 16 USD Discount Coupon: **NSZUBG3X**

**Email:** [support@examsempire.com](mailto:support@examsempire.com)

**Check our Customer Testimonials and ratings  
available on every product page.**

**Visit our website.**

**<https://examsempire.com/>**