

# Healthcare BC-ADM

**AADE Diabetes Management - Advanced (BC-ADM)**

**For More Information – Visit link below:**

**<https://www.examsempire.com/>**

**Product Version**

- 1. Up to Date products, reliable and verified.**
- 2. Questions and Answers in PDF Format.**



**<https://examsempire.com/>**

**Visit us at: <https://www.examsempire.com/bc-adm>**

# Latest Version: 6.0

## Question: 1

A child is diagnosed with type I diabetes. What would be normal?

- A. Regression.
- B. Advanced development.
- C. Normal development.
- D. All of the above.

**Answer: A**

Explanation:

Understanding the effects of a chronic illness such as type I diabetes on a child involves considering multiple dimensions of a child's life, including their developmental progression. Normally, children follow a predictable path of developmental milestones, but the onset of a chronic condition can temporarily disrupt this progression.

Regression in development is a common response among children who experience significant changes in their health status. This regression can manifest as a temporary reversion to earlier behaviors or skills. For instance, a child who has been toilet trained might begin experiencing accidents, or a previously independent child might become more clingy or dependent on adults. This kind of regression is a normal psychological reaction to the stress and uncertainty brought about by illness, as the child seeks comfort in more familiar and secure behaviors.

While regression is a normal response, it is crucial for caregivers and healthcare providers to monitor these changes closely. The goal is to support the child through this period without letting the regression hinder their overall developmental trajectory. Interventions might include providing additional emotional support, maintaining a routine as much as possible, and involving the child in their care regimen to foster a sense of control.

In the context of the question regarding what would be normal for a child diagnosed with type I diabetes, the answer "Regression" acknowledges that it is common for children to show some temporary developmental backsliding as they adjust to the diagnosis and management of their condition. This is considered within the normal spectrum of responses and does not necessarily indicate a problem, provided that the child receives appropriate support to return to their typical developmental path eventually.

Overall, understanding and supporting a child with type I diabetes requires a sensitive approach that considers both their physical and psychological needs. Recognizing and addressing regression as a normal response can help in devising appropriate strategies to support the child's well-being and development during this challenging time.

## Question: 2

Which change in digestion typically occurs in patients as they age?

- A. Increased saliva production.
- B. Decreased saliva production.
- C. Increased thirst.
- D. Faster transfer time.

**Answer: B**

Explanation:

As individuals age, several physiological changes occur that impact various systems in the body, including the digestive system. One common change observed in the elderly is decreased saliva production. Saliva plays a critical role in the initial stages of digestion by moistening food and beginning the process of chemical digestion through enzymes such as amylase. Decreased saliva production can lead to difficulties in chewing and swallowing food, which is medically referred to as dysphagia. Decreased saliva production in older adults can result from multiple factors. These include systemic diseases like diabetes and Sjögren's syndrome, which directly impair the function of the salivary glands. Additionally, many medications commonly prescribed for older adults, such as antihistamines, diuretics, and medications for hypertension, can also reduce saliva output as a side effect. Furthermore, diminished oral health in elderly populations, often aggravated by less effective dental care, can exacerbate the problems associated with reduced saliva production. Poor oral health can lead to a further decrease in the functional capacity of salivary glands, compounding the challenges of food ingestion and digestion.

The reduction in saliva production is not just a matter of discomfort. It has broader implications for nutrition and health in elderly individuals. Adequate saliva is necessary not only for the digestion of carbohydrates but also for the perception of taste. A decrease in saliva can lead to diminished taste sensitivity, which may decrease appetite and possibly lead to nutritional deficiencies. Additionally, saliva has antibacterial properties that help maintain oral hygiene. A decrease in saliva production can therefore increase the risk of oral infections and systemic infections stemming from the mouth. In summary, decreased saliva production is a significant change in the digestive process of aging patients, impacting not only their ability to process food but also their overall oral health and nutritional status. This change underscores the importance of monitoring and managing oral health and dietary habits in elderly individuals to ensure they maintain adequate nutrition and quality of life.

### Question: 3

Which of the following screening tests will correlate with a diabetic foot infection?

- A. Temperature.
- B. Sweats.
- C. Leukocytosis.
- D. Erythrocyte sedimentation rate.

**Answer: D**

Explanation:

When considering the appropriate screening tests for detecting a diabetic foot infection, various indicators are evaluated, but not all may have a strong correlation with an infection in patients with

diabetes. A diabetic foot infection is a common complication in individuals with diabetes, which can lead to serious outcomes if not promptly diagnosed and treated.

Temperature measurement alone may not be a reliable indicator. While elevated temperature can suggest an infection, diabetic individuals, especially those with neuropathy, might not exhibit typical signs of infection such as fever. This can be due to altered inflammatory responses or less perceivable symptoms because of nerve damage.

Leukocytosis, which refers to an increased number of white blood cells, might seem like a straightforward indicator of infection. However, in the context of diabetic foot infections, its presence is inconsistent. Many patients with a diabetic foot infection may not exhibit leukocytosis, making it an unreliable sole indicator for infection.

The Erythrocyte Sedimentation Rate (ESR) is considered a useful marker in the context of diabetic foot infections. It is a non-specific test that measures the rate at which red blood cells sediment in a period of one hour. An elevated ESR is indicative of an inflammatory response, which is typical in infections. In the majority of cases involving diabetic foot infections, the ESR is elevated, reflecting the body's response to the infection. This makes ESR a more reliable marker compared to temperature and leukocyte count in the context of diabetic foot infections.

Therefore, among the options provided, the Erythrocyte Sedimentation Rate (ESR) is the correct answer as it is more likely to be elevated in patients with a diabetic foot infection, providing a useful correlate for the presence of an infection. It is important for healthcare providers to consider multiple factors and possibly a combination of tests when diagnosing diabetic foot infections to ensure accurate and timely management.

### Question: 4

Which of the following is a step to consider when modifying a diabetic care plan?

- A. Add new self management goals.
- B. Examine the intervention.
- C. Re-evaluate the patient and their care.
- D. All of the above.

**Answer: D**

Explanation:

When considering modifications to a diabetic care plan, it's crucial to adopt a comprehensive approach that addresses various aspects of patient management. The correct answer to the question is "All of the above," as each option listed plays a vital role in refining and adapting the care plan to better meet the patient's needs. Here's a breakdown and expansion of each step mentioned:

**\*\*Gathering data on the diagnosis\*\*:** This initial step involves reviewing the current health status and medical history of the patient. It is crucial to ensure that the diagnosis is accurate and to assess any new symptoms or changes that might affect the patient's condition. This data serves as the foundation for any adjustments in the care plan.

**\*\*Setting new priorities\*\*:** Diabetes management can involve multiple health concerns, such as blood sugar control, diet, exercise, and managing co-existing conditions like hypertension or dyslipidemia. Prioritizing these aspects based on the patient's current health status helps in focusing on the most critical areas first.

**\*\*Changing the diagnosis if necessary\*\***: If new data suggests that the initial diagnosis was incomplete or incorrect, it might be necessary to adjust the diagnosis. This could involve identifying complications or secondary conditions that need to be addressed.

**\*\*Examining the diagnosis\*\***: A continuous re-evaluation of the diagnosis ensures that it remains relevant as the patient's condition evolves. This might include regular monitoring of blood glucose levels, HbA1c tests, and other relevant diagnostics.

**\*\*Changing the goals that are unattainable\*\***: Some goals set at the beginning of the care plan might become unrealistic as the situation changes. It's important to adjust these goals to ensure they remain achievable, which helps in maintaining patient motivation and compliance.

**\*\*Adding new goals\*\***: As the patient's condition progresses or if new health issues arise, setting new goals becomes necessary. These should be specific, measurable, attainable, relevant, and time-bound (SMART goals).

**\*\*Getting rid of goals that will not be met\*\***: This involves removing or revising goals that are no longer relevant or feasible, which could be due to changes in the patient's health status or life circumstances.

**\*\*Examining the intervention\*\***: This step involves assessing the current treatment plan, including medication, diet, exercise, and other management strategies to determine their effectiveness and suitability.

**\*\*Changing intervention processes\*\***: If current interventions are not yielding the expected outcomes, changes may be required. This could mean altering medications, introducing new therapies, or adjusting dietary and exercise recommendations.

**\*\*Incorporating new factors for goal achievement\*\***: Consideration of new or previously unaddressed factors such as mental health, social support, and access to healthcare resources can enhance the effectiveness of the care plan.

**\*\*Re-evaluating the patient\*\***: Regular follow-ups and patient assessments are essential to gauge progress, understand patient concerns, and make further adjustments to the care plan. This holistic view ensures that the care plan remains aligned with the patient's health needs and lifestyle.

In summary, modifying a diabetic care plan is a dynamic and multifaceted process requiring ongoing evaluation and adaptation. Every aspect, from diagnosis to intervention and goal-setting, must be regularly scrutinized and adjusted to ensure optimal patient care and health outcomes. The answer "All of the above" encapsulates the necessity of considering all these steps in the modification of a diabetic care plan.

## Question: 5

A serving of carbohydrates is \_\_\_\_\_.

- A. 12g.
- B. 10g.
- C. 17g.
- D. 15g.

**Answer: D**

Explanation:

Understanding the correct serving sizes of carbohydrates is crucial for individuals, particularly those managing conditions like diabetes or metabolic syndrome. The standard measure for a carbohydrate

serving is 15 grams. This unit of measure helps in quantifying the intake of carbohydrates in a consistent and controlled manner, facilitating better management of blood glucose and lipid levels in the body. It is essential that patients learn how to accurately measure and understand what constitutes a 15g serving of carbohydrates. This could include foods like a small piece of fruit, a slice of bread, or a third of a cup of cooked rice. Being precise with these measurements allows individuals to maintain a balanced diet and avoid the spikes in blood sugar levels that can occur if too many carbohydrates are consumed in one sitting.

Moreover, it is not just about the quantity but also about the distribution of carbohydrate intake throughout the day. Spreading carbohydrate consumption evenly across meals and snacks helps in maintaining steady blood glucose levels and prevents the highs and lows that can be detrimental to health. This is particularly important for those with insulin sensitivities or those who are on medications that affect insulin levels.

In conclusion, a 15g serving of carbohydrates is a foundational concept in dietary planning for glucose and lipid control. Educating patients about this helps them make informed choices about their diets, contributing to better overall health management. This understanding empowers individuals to take active control of their dietary habits, leading to improved health outcomes and quality of life.

### Question: 6

An elderly patient with type II diabetes keeps developing dehydration. How much water should the patient be encouraged to drink each day?

- A. 24 oz.
- B. 60 oz.
- C. 32 oz.
- D. 48 oz.

**Answer: D**

Explanation:

Dehydration is a common concern for elderly patients, especially those with type II diabetes. In such patients, dehydration can exacerbate health issues by causing an increase in blood glucose levels and the concentration of administered medications. This can lead to complications such as hyperglycemia or even more severe effects due to altered drug efficacy and metabolism.

To prevent these complications, it is essential for diabetic patients to maintain adequate hydration. For an elderly patient with type II diabetes who is prone to dehydration, it is generally recommended that they consume at least 48 ounces of water each day. This amount helps in maintaining proper hydration and supports the kidney's ability to manage blood glucose levels and eliminate waste products efficiently.

However, it is crucial to note that while 48 ounces is a general recommendation, the specific daily water intake might need to be adjusted based on individual health conditions and under medical advice. For instance, patients who have conditions like congestive heart failure or renal issues may require a different intake level to avoid fluid overload, which could exacerbate their condition.

In summary, ensuring that an elderly patient with type II diabetes drinks around 48 ounces of water daily is a foundational step in managing their condition. Yet, this should always be tailored to the patient's overall health profile and other existing medical conditions, following the guidance of healthcare professionals.

### Question: 7

A patient with symptoms of hypoglycemia tests her blood glucose 15 minutes after eating 15 grams of simple carbohydrates. Her blood glucose is 60 mg/dL. What should she do next?

- A. Consume a small meal.
- B. Wait 15 minutes and check her blood sugar again.
- C. Consume 15 grams of protein.
- D. Consume an additional 15-20 grams of simple carbohydrates or glucose.

**Answer: D**

Explanation:

In the scenario described, the patient has already consumed 15 grams of simple carbohydrates in response to symptoms of hypoglycemia. Despite this initial intake, her blood glucose level remains low at 60 mg/dL, which is below the normal threshold. According to standard guidelines for managing hypoglycemia, the advisable next step is to consume an additional 15-20 grams of simple carbohydrates or glucose.

This recommendation is based on the "15-15 Rule," which suggests that if blood glucose is still below 70 mg/dL after the first 15 grams of carbohydrates, another 15-20 grams should be consumed. This helps to ensure that the blood sugar levels rise to a safer range. The approach involves a repeated cycle of consuming carbohydrates, waiting approximately 15 minutes, and then re-checking blood glucose levels. Simple carbohydrates are recommended because they are absorbed quickly by the body, raising blood sugar levels more rapidly than complex carbohydrates or proteins. Foods that typically contain about 15 grams of simple carbohydrates include glucose tablets, honey, regular soda, or fruit juice. After re-administering the carbohydrates, it's crucial for the patient to retest her blood glucose after another 15 minutes to ensure that it has returned to a safer level.

If, after the second round of carbohydrate intake, the blood sugar remains below 70 mg/dL, this cycle may need to be repeated, or medical assistance might be required. It is important for individuals with frequent episodes of hypoglycemia, or those who are on medication such as insulin, to discuss their management plan with a healthcare provider to tailor it based on their specific needs and experiences. Additionally, once blood glucose levels are stabilized, consuming a small meal or snack that includes protein and complex carbohydrates can help sustain blood glucose levels and prevent further drops.

### Question: 8

A clear, concise statement of the goal of a research study is known as which of the following?

- A. Hypothesis.
- B. Research problem.
- C. Research purpose.
- D. Research topic.

**Answer: C**

Explanation:

The term "Research purpose" refers to a clear, concise statement that outlines the primary goal or objective of a research study. This statement is crucial as it guides the entire research process, helping researchers maintain focus and clarity about what they intend to achieve. The research purpose is fundamentally about defining what the researcher aims to discover, examine, or establish through their study.

In contrast to other elements of a research project such as the hypothesis, research problem, or research topic, the research purpose serves a unique role. While a hypothesis is a specific, testable prediction about what you expect to happen in your study, the research purpose is broader and provides the overarching reason for why the study is being conducted. The research problem, on the other hand, defines the issue that the research is intended to address, often pointing out gaps in knowledge that the research aims to fill. The research topic is generally the subject area or the broad field within which the research is conducted.

The significance of having a well-defined research purpose cannot be overstated. It not only directs the research methodology and influences the choice of tools and techniques but also assists in framing the research questions and/or hypotheses. A clearly articulated purpose aids in keeping the study aligned with its initial goals, preventing scope creep, and ensuring that the outcomes are relevant and valuable. In academic or professional contexts, the research purpose is typically presented at the beginning of a research proposal or report. It sets the stage for introducing the detailed elements of the research, such as the methodology, literature review, data analysis, and findings. For researchers, being able to articulate a clear research purpose is indicative of a thorough understanding of their field and the specific challenges or questions they are addressing.

Thus, when crafting a research purpose, it is essential for researchers to be succinct yet detailed enough to convey the essence of what the study seeks to achieve. This involves not only stating the intent of the study but also highlighting its significance and potential impact on the field or society. A well-defined research purpose is a hallmark of a thoughtfully planned and potentially impactful study.

### Question: 9

All but which of the following have been shown to increase a person's risk of Type 2 diabetes?

- A. Alcohol use.
- B. Sedentary lifestyle.
- C. A diet which incorporates red meat.
- D. Stress.

**Answer: C**

Explanation:

It appears there was some confusion in the structuring of your question and answer. Let's clarify the question and provide a more detailed explanation for the correct answer.

Question

A diet which incorporates red meat 3. Sedentary lifestyle 4. Stress

Correct Answer: A diet which incorporates red meat

Explanation: Type 2 diabetes is a metabolic disorder that is primarily characterized by high blood glucose in the context of insulin resistance and relative insulin deficiency. Several lifestyle factors have been



identified that increase the risk of developing Type 2 diabetes. These include excessive alcohol use, maintaining a sedentary lifestyle, and experiencing high levels of stress.

Alcohol use can lead to chronic pancreatitis, which impairs the pancreas' ability to secrete insulin, thereby directly contributing to diabetes. Furthermore, heavy drinking can lead to weight gain and changes in metabolism, both of which are risk factors for diabetes.

Sedentary lifestyles contribute to the development of Type 2 diabetes by promoting obesity, decreasing insulin sensitivity, and reducing the muscle mass that is crucial for proper glucose metabolism.

Stress affects blood sugar levels directly by increasing the release of glucagon and cortisol, which in turn increase blood glucose levels. Chronic stress can also lead to poor lifestyle choices such as unhealthy eating and reduced physical activity, which further increase diabetes risk.

On the other hand, the consumption of red meat, while often cited as a risk factor for various health conditions, shows a more complex relationship with Type 2 diabetes. Studies suggest that while excessive consumption of processed and unprocessed red meat may be associated with an increased risk of diabetes, moderate consumption within a balanced diet does not independently raise diabetes risk significantly. It is the mode of preparation, the type of meat, and the overall dietary pattern that play more pivotal roles in the impact of red meat consumption on diabetes risk.

In conclusion, among the options given, a diet which incorporates red meat, if consumed in moderation and part of a balanced diet, does not necessarily increase the risk of Type 2 diabetes to the same extent as the other listed factors such as alcohol use, a sedentary lifestyle, and high stress levels.

### Question: 10

Unresolved conflict among members of a healthcare team can have all but which of the following effects?

- A. It can jeopardize patient safety.
- B. It can compromise the quality of patient care
- C. It can increase the quality of patient care.
- D. It can negatively impact the patient's trust in the healthcare team.

**Answer: C**

Explanation:

Unresolved conflicts among members of a healthcare team can lead to several negative outcomes, but not an increase in the quality of patient care. When team members are in conflict, communication often breaks down. Effective communication is critical in healthcare settings where the stakes are high, and information needs to be clear and precise. Miscommunications or withholding of information can lead to errors in patient care, potentially jeopardizing patient safety.

Furthermore, unresolved conflicts can create a stressful work environment, which can affect the mental and emotional well-being of healthcare providers. This stress can reduce their ability to focus and perform their duties efficiently, which can compromise the quality of care that patients receive. In high-stress situations, the likelihood of making mistakes increases, and in a healthcare setting, this can have serious consequences for patient outcomes.

Another significant impact of unresolved conflicts is the degradation of team cohesion. Healthcare relies heavily on the teamwork of various professionals who must collaborate to provide the best care. When team members are in conflict, the collaboration necessary for patient care can suffer. This lack of

cohesion can lead to disjointed patient care where important aspects of a patient's treatment might be overlooked or mishandled.

It is also important to consider the patient's perspective in situations where there is evident discord among the healthcare team. Patients who perceive conflict may experience decreased trust and confidence in their care providers. This perception can lead to decreased patient satisfaction and could deter patients from following medical advice or returning for necessary follow-up care.

In conclusion, unresolved conflict among healthcare team members does not lead to an increase in the quality of patient care. Instead, it compromises communication, increases stress levels, diminishes team cohesion, and can lead to a loss of patient trust—all of which negatively impact the quality of care provided to patients.

**Thank You for Trying Our Product**

**Special 16 USD Discount Coupon: NSZUBG3X**

**Email:** [support@examsempire.com](mailto:support@examsempire.com)

**Check our Customer Testimonials and ratings  
available on every product page.**

**Visit our website.**

**<https://examsempire.com/>**