

# Canadian CNCC

**Canadian Critical Care Nursing Examination (CNCC)**

**For More Information – Visit link below:**

**<https://www.examsempire.com/>**

**Product Version**

- 1. Up to Date products, reliable and verified.**
- 2. Questions and Answers in PDF Format.**



**<https://examsempire.com/>**

**Visit us at: <https://www.examsempire.com/cncc>**

# Latest Version: 6.0

## Question: 1

The outermost layer of the stomach and intestinal wall is which of the following?

- A. serosa
- B. mucosa
- C. submucosa
- D. smooth muscle layer

**Answer: A**

Explanation:

The correct answer to the question regarding the outermost layer of the stomach and intestinal wall is "serosa." The gastrointestinal tract, specifically the stomach and intestines, is composed of several layers that each play important roles in digestion and the structural integrity of these organs.

Starting from the inside, the first layer is the **mucosa**. This is the innermost lining that comes into direct contact with the contents of the gastrointestinal tract. It is responsible for most digestive, absorptive, and secretory functions. The mucosa itself is made up of three sub-layers: the epithelium, the lamina propria, and the muscularis mucosae.

Next is the **submucosa**. This layer lies just outside the mucosa and consists of dense and loose connective tissue that supports the mucosa. It contains blood vessels, nerves, and glands that are crucial for the transport of nutrients and the secretion of digestive enzymes.

Following the submucosa are the **muscular layers**. Typically, there are two main muscle layers – the inner circular and the outer longitudinal layers. These layers are made of smooth muscle fibers that facilitate peristalsis, the rhythmic contractions that propel food through the digestive tract.

The outermost layer of the stomach and intestinal wall is the **serosa**. This layer is a smooth, slippery tissue derived from the visceral peritoneum; it covers the external surface of most of the gastrointestinal tract. The serosa serves primarily as a protective layer and helps to reduce friction between the stomach and intestines and other abdominal organs. The serosa secretes a serous fluid that lubricates and allows smooth movement of the gastrointestinal tract within the abdominal cavity. Understanding the structure of these layers is crucial for comprehending how the stomach and intestines function, how they interact with other systems in the body, and how diseases affecting these layers can impact overall health. The serosa, as the outermost layer, plays a crucial role in both protection and structural integrity.

## Question: 2

Decreased platelet production can result in which of the following?

- A. disseminated intravascular coagulation
- B. anemia
- C. immunosuppression

D. thrombocytopenia

**Answer: D**

Explanation:

Decreased platelet production can result in thrombocytopenia, a medical condition characterized by a lower than normal count of platelets in the blood. Platelets, or thrombocytes, are small blood cell fragments crucial for normal blood clotting. When their numbers fall below a certain threshold, the ability of the blood to clot is significantly impaired, leading to increased risks of bleeding.

Thrombocytopenia can be caused by various factors that interfere with the production of platelets in the bone marrow. Some of these factors include bone marrow disorders such as leukemia or myelodysplastic syndromes, certain infections, specific medications, chemotherapy drugs, and radiation therapy. Additionally, heavy alcohol consumption and exposure to toxic chemicals can also suppress platelet production.

The clinical manifestations of thrombocytopenia depend on the severity of the condition. Mild cases might not present any symptoms and may only be discovered during routine blood tests. However, as the condition progresses, symptoms can include frequent and easy bruising, prolonged bleeding from cuts, spontaneous bleeding from the gums or nose, excessively heavy menstrual flows, and petechiae (small red spots on the skin caused by bleeding under the skin). Severe thrombocytopenia can lead to more serious bleeding episodes, such as gastrointestinal bleeding or intracerebral hemorrhage, which can be life-threatening.

Diagnosis of thrombocytopenia involves a complete blood count (CBC) test, which will show a reduced number of platelets. Further diagnostic tests like a bone marrow biopsy might be necessary to determine the underlying cause of the decreased platelet production. Treatment of thrombocytopenia is directed towards the underlying cause and might include medications to boost platelet production, blood or platelet transfusions, and treatments specific to any underlying disease.

It is important to address thrombocytopenia promptly to manage symptoms effectively and prevent complications such as severe bleeding. Regular monitoring and medical check-ups are essential for patients diagnosed with this condition to keep track of their platelet counts and ensure appropriate interventions are made when necessary.

### Question: 3

Which of the following statements about intracranial pressure (ICP) is least accurate?

- A. ICP is the pressure exerted by the intracranial contents, brain, tissue, blood, and cerebrospinal fluid within the cranium.
- B. The ventricular catheter with external strain gauge transducer is considered the gold standard for ICP monitoring.
- C. Increased ICP occurs when the brain's ability to compensate for increased volume exceeds the intracranial volume.
- D. Increased ICP contributes to secondary neuronal injury.

**Answer: C**

Explanation:

Intracranial pressure (ICP) is defined as the pressure within the skull, exerted by components such as brain tissue, blood, and cerebrospinal fluid (CSF). These components must maintain a delicate balance to ensure proper brain function and to prevent damage due to excessive pressure.

The Monro-Kellie doctrine explains that the cranial compartment is incompressible, and the volume inside the cranium is a fixed sum of the brain tissue, blood, and CSF. Any increase in one of these components must be compensated by a decrease in one or more of the other components to maintain normal ICP. When the capacity for compensation is exceeded, ICP increases.

The statement that "Increased ICP occurs when the brain's ability to compensate for increased volume exceeds the intracranial volume" is incorrect and thus identified as the least accurate. The phrasing suggests that it is the ability to compensate that is being exceeded, which is misleading. In reality, increased ICP occurs when the intracranial volume increases beyond the brain's ability to compensate, typically due to swelling, hemorrhage, or other pathologies that add volume to the intracranial contents. To monitor ICP, various methods are employed, among which the ventricular catheter with an external strain gauge transducer is considered the gold standard. This method involves placing a catheter in one of the brain's ventricles, allowing direct measurement of pressure and facilitating the drainage of CSF to temporarily reduce ICP if necessary.

It is crucial to manage elevated ICP effectively because prolonged increased pressure can lead to serious consequences, such as brain herniation, impaired blood flow to the brain, and ultimately, secondary neuronal injury. These secondary injuries can significantly worsen outcomes for patients with conditions like traumatic brain injury, stroke, or brain tumors. Managing ICP involves medical and surgical interventions aimed at reducing the volume within the skull and controlling factors that contribute to raised ICP.

### Question: 4

The life-threatening augmentation of the signs and symptoms of hyperthyroidism is known as which of the following?

- A. syndrome of inappropriate antidiuretic hormone secretion
- B. thyrotoxicosis
- C. diabetes insipidus
- D. myxedema coma

**Answer: B**

Explanation:

Thyrotoxicosis, also known as a thyroid storm, is a severe and potentially life-threatening health condition that arises from excessively high levels of thyroid hormones (thyroxine or T4 and triiodothyronine or T3), exacerbating the symptoms of hyperthyroidism. This condition represents an extreme state of hyperthyroidism and is considered a medical emergency.

Thyrotoxicosis can be precipitated by various factors such as infections, trauma, surgery, uncontrolled diabetes, pregnancy, or discontinuation of thyroid medication in hyperthyroid patients. The excessive levels of thyroid hormones can lead to a rapid increase in metabolic rate, causing symptoms such as fever, sweating, palpitations, arrhythmias, nausea, vomiting, diarrhea, severe weakness, and even heart failure.

The diagnosis of thyrotoxicosis is primarily based on clinical symptoms and is confirmed through biochemical tests showing elevated levels of thyroid hormones and suppressed thyroid-stimulating

hormone (TSH) levels. Immediate and aggressive treatment is crucial to manage the symptoms and prevent complications. Treatment options usually include antithyroid medications, beta-blockers to control heart rate, steroids, and in some cases, emergency surgery.

Although thyrotoxicosis is a rare complication of hyperthyroidism, it is critical to recognize and treat it promptly. Most cases of hyperthyroidism are well-managed with antithyroid drugs, which helps in significantly reducing the risk of developing thyrotoxicosis. However, it remains a critical condition that requires immediate medical intervention to avoid severe outcomes or death.

### Question: 5

Which of the following statements about consent for clinical care is least accurate?

- A. Expressed consent is given directly by written or verbal words.
- B. Implied consent is presumed in emergency situations.
- C. Partial consent is not an option.
- D. Consent can be implied by a patient's behavior.

**Answer: C**

Explanation:

Understanding the concept of consent within clinical care is essential for both healthcare providers and patients. Consent can be categorized in various forms, including expressed, implied, and partial consent. Each type has its specific conditions and implications.

Expressed consent is when consent is clearly and explicitly communicated by the patient, either verbally or in writing. This type of consent leaves little to no ambiguity about the patient's wishes regarding specific medical treatments or procedures. For instance, a patient signing a consent form for a surgical procedure is an example of expressed written consent.

Implied consent, on the other hand, is not given directly but rather inferred from a person's actions or the context of a situation. Commonly recognized in emergency scenarios, implied consent assumes that a patient would agree to necessary medical interventions if they were able to provide consent. For example, an unconscious patient brought into an emergency department is treated under the assumption of implied consent, as immediate care is necessary to preserve life or prevent serious harm.

Partial consent is where there appears to be some confusion or misunderstanding. Contrary to the statement that "Partial consent is not an option," partial consent is indeed a valid and frequently utilized form of consent in clinical settings. Partial consent occurs when a patient agrees to certain parts of a treatment plan but declines others. For example, a patient may consent to undergo diagnostic tests but decline any form of surgical intervention that might be suggested following the diagnosis. This demonstrates that the patient's autonomy is respected to choose which medical interventions they are comfortable with and which they are not.

The least accurate statement in the context provided is, therefore, "Partial consent is not an option." In clinical practice, it is crucial for healthcare providers to recognize and respect partial consent as it upholds the patient's right to make informed decisions about their own health care. Misunderstanding this can lead to ethical and legal issues, potentially compromising patient trust and adherence to recommended medical procedures.

In summary, understanding the different types of consent is fundamental in ensuring ethical medical practice and patient autonomy. Each type of consent serves a specific purpose and is applicable under

different circumstances, with partial consent playing a critical role in personalized and patient-centered care.

### Question: 6

Abraham Maslow characterized needs in terms of a hierarchy. The ultimate goal in this hierarchy is which of the following?

- A. self-esteem
- B. self-actualization
- C. love and a sense of belonging
- D. safety and security

**Answer: B**

Explanation:

Abraham Maslow, a renowned psychologist, developed a theory in psychology that outlines a hierarchy of needs. This hierarchy is often depicted as a pyramid divided into several levels, with the most basic needs at the bottom and the highest level of needs at the top. Maslow's Hierarchy of Needs is structured as follows: physiological needs (such as food and water), safety needs, love and belonging needs, esteem needs, and finally, at the top, self-actualization needs.

According to Maslow, self-actualization represents the ultimate goal in this hierarchical model. It is the realization of an individual's potential, self-fulfillment, seeking personal growth, and peak experiences. Maslow described self-actualization as a person's need to be and do that which the person was "born to do." Essentially, it is about becoming everything that one is capable of becoming, which may manifest as pursuing talents, capabilities, and creativity.

It is crucial to understand that Maslow's theory posits that lower-level needs must be satisfied before the higher-level needs can be addressed. For instance, if an individual is struggling with basic physiological needs such as hunger and thirst, their motivation to address higher needs like self-esteem or self-actualization will be suppressed until those basic needs are met. This progression through the hierarchy explains why self-actualization sits at the apex of the pyramid: it is pursued only after all other lower, more fundamental needs are adequately fulfilled.

In practical terms, self-actualization might manifest in various forms, including artistic expression, pursuit of knowledge, involvement in sports, or commitment to community service, depending on an individual's personal interests and capacities. Not everyone will reach the stage of self-actualization, and the journey toward this stage can often be influenced by an individual's circumstances, including critical illnesses or other significant life challenges, which may necessitate a refocus on more basic needs such as physiological requirements and safety.

Thus, when asked about the highest goal in Maslow's hierarchy, the answer is self-actualization. This stage is not about basic survival or social belonging but achieving one's fullest potential and realizing one's deepest talents and capabilities.

### Question: 7

When observing a patient's musculoskeletal development, if the sternum protrudes instead of being lower than the adjacent hemithoraces this is known as which of the following?

- A. pectus excavatum
- B. pectus carinatum
- C. scoliosis
- D. kyphosis

**Answer: B**

Explanation:

The correct answer to the question is pectus carinatum.

Pectus carinatum, also known as pigeon chest, is a deformity of the chest characterized by a protrusion of the sternum and ribs. This protrusion occurs when the cartilage that connects the ribs to the breastbone grows excessively, causing the sternum to jut outward rather than lying flat or being slightly concave. The visual effect is that the chest appears to be pushed outward, which contrasts with the normal or slightly concave alignment of the sternum relative to the adjacent hemithoraces.

In contrast, pectus excavatum, commonly referred to as sunken chest, is another condition involving the sternum but is characterized by a concave appearance. In this condition, the sternum is depressed inward toward the spine, creating a sunken appearance in the chest. This is the opposite of the protrusion seen in pectus carinatum.

Scoliosis and kyphosis, the other options listed, are different types of spinal deformities and do not involve the sternum directly. Scoliosis is characterized by a lateral (sideways) curvature of the spine, whereas kyphosis involves an exaggerated forward rounding of the back. Both of these conditions affect the spine and not primarily the chest structure as in the cases of pectus carinatum and pectus excavatum.

Therefore, when observing a patient whose sternum protrudes outward, the correct term for this musculoskeletal development is pectus carinatum. This condition can vary in severity and may cause physical discomfort and psychological issues due to its appearance, but it is distinguishable from other chest and spinal deformities by its specific characteristic of sternum protrusion.

### Question: 8

In terms of the pathophysiology of coronary artery disease which of the following statements is accurate?

- A. Plaque in the coronary arteries is a pearly white accumulation in the intimal lining of the artery.
- B. Fatty fibrous plaque in an artery can rupture and form either a mural thrombus or an occlusive thrombus.
- C. Atherosclerotic process causes decreases in blood flow and oxygen supply to the myocardium.
- D. all of the above

**Answer: D**

Explanation:

Coronary artery disease (CAD) is a significant medical condition characterized by the narrowing or blockage of the coronary arteries, primarily due to the process of atherosclerosis. Atherosclerosis is a chronic inflammatory response in the walls of arteries, in large part due to the accumulation of

macrophage white blood cells and promoted by low-density lipoproteins (plasma proteins that carry cholesterol and triglycerides). This pathophysiological process leads to the formation of plaques, which are not pearly white but typically consist of a fatty, fibrous core covered by a fibrous cap.

These plaques are what accumulate in the intimal lining of the artery. Initially, these plaques might not significantly affect blood flow but as they grow larger, they can restrict blood flow. Furthermore, if the fibrous cap over a plaque ruptures, it can lead to the formation of a blood clot (thrombus). This thrombus can either partially or completely block the blood flow in the coronary artery. Depending on the extent and location of the blockage, this can lead to symptoms such as angina (chest pain) or more severe outcomes like myocardial infarction (heart attack).

The occlusion of coronary arteries through these mechanisms leads to decreased blood flow and a reduced oxygen supply to the myocardium (heart muscle). This ischemia (reduced blood flow) and hypoxia (reduced oxygen supply) can impair heart muscle function, which in severe cases, can lead to heart failure or death.

The statement in the question regarding plaque being a "pearly white accumulation" is inaccurate.

Plaque within the arteries is typically yellowish and consists of lipids, calcium, and other cellular waste products, not a pearly white substance. This description might be a simplification or misconception. The rest of the statements provided correctly describe the progressive nature of CAD, the role of atherosclerosis, and the potential acute complications arising from plaque rupture and thrombosis.

In summary, CAD is a complex and progressive disease involving the build-up of atherosclerotic plaques in the coronary arteries, which can lead to partial or complete occlusion of these arteries. The consequences of such occlusions can be severe, affecting the heart's ability to function efficiently and increasing the risk of heart attacks and heart failure. Accurate understanding and description of these processes are crucial for proper diagnosis, management, and treatment of coronary artery disease.

## Question: 9

Peritoneal dialysis works on which of the following principles?

- A. diffusion
- B. osmosis
- C. both diffusion and osmosis
- D. none of the above

**Answer: C**

Explanation:

Peritoneal dialysis is a medical process used to remove waste products and excess fluid from the blood when the kidneys are no longer able to perform these functions naturally. This form of dialysis utilizes the patient's peritoneum in the abdomen as a membrane across which fluids and dissolved substances (electrolytes, urea, glucose, albumin and other small molecules) are exchanged from the blood.

The effectiveness of peritoneal dialysis hinges on the principles of both diffusion and osmosis, which are two types of passive transport mechanisms involving the movement of molecules across a semipermeable membrane.

**\*\*Diffusion\*\*** is the process by which molecules move from an area of higher concentration to an area of lower concentration. In the context of peritoneal dialysis, this means that waste products like urea and creatinine, which are present in high concentrations in the blood but lower in the dialysis fluid



(dialysate), move across the peritoneal membrane into the dialysate. Similarly, excess electrolytes and other solutes also diffuse down their concentration gradients from the blood into the dialysate.

**\*\*Osmosis\*\***, on the other hand, involves the movement of water (solvent) across a semipermeable membrane from an area of lower solute concentration to an area of higher solute concentration. In peritoneal dialysis, the dialysate typically contains a sugar solution, often dextrose, which is at a higher concentration than the fluid inside the blood vessels. This creates an osmotic gradient that causes water to move from the blood into the dialysate, thereby helping to remove excess fluid from the body. Together, these processes of diffusion and osmosis in peritoneal dialysis help to regulate the chemical composition of the blood, adjusting levels of fluid, electrolytes, and waste materials, effectively mimicking the filtration function of healthy kidneys. The dialysate is periodically introduced into and removed from the peritoneal cavity, ensuring continuous treatment and maintenance of safe, balanced blood chemistry.

### Question: 10

Blood in the pleural space of a trauma patient is known as which of the following?

- A. hemothorax
- B. open pneumothorax
- C. pulmonary contusion
- D. blunt cardiac injury

**Answer: A**

Explanation:

The correct answer to the question "Blood in the pleural space of a trauma patient is known as which of the following?" is hemothorax. A hemothorax occurs when blood accumulates in the pleural cavity, which is the space between the lungs and the chest wall. This condition is commonly caused by chest trauma which may result from accidents, injuries, or surgical complications.

The presence of blood in the pleural space can compromise lung function by limiting the expansion of the lungs during breathing. This can lead to respiratory distress and inadequate oxygenation of the blood. The severity of a hemothorax can vary depending on the amount of blood that has accumulated. In severe cases, it can be a life-threatening condition that requires immediate medical intervention.

One indicator of the potential severity of a hemothorax is the involvement of the first and second ribs.

These ribs are located high in the rib cage and are more protected than the lower ribs. As a result, fractures to these ribs usually indicate a high-impact force. Such force can also damage major blood vessels or the heart, which are situated near these ribs, thereby increasing the risk of a hemothorax.

In the context of trauma, other related conditions that might be considered include open pneumothorax, pulmonary contusion, and blunt cardiac injury. An open pneumothorax involves an open wound to the chest that allows air to enter the pleural space, collapsing the lung. A pulmonary contusion refers to bruising of the lung tissue, which can lead to swelling and bleeding within the lung. Blunt cardiac injury describes trauma to the heart itself, which can affect its function and potentially lead to other complications like cardiac tamponade or arrhythmias.

In summary, a hemothorax is characterized by the accumulation of blood in the pleural space, typically resulting from traumatic injury. It is a serious medical condition that necessitates prompt and effective treatment to prevent complications such as hypoxia and shock. Management may include draining the blood from the pleural space and addressing any underlying injuries to prevent further bleeding.

**Thank You for Trying Our Product**

**Special 16 USD Discount Coupon: NSZUBG3X**

**Email:** [support@examsempire.com](mailto:support@examsempire.com)

**Check our Customer Testimonials and ratings  
available on every product page.**

**Visit our website.**

**<https://examsempire.com/>**