

# Dental NDAEB

**NDAEB Canadian Certified Dental Assistant**

**For More Information – Visit link below:**

**<https://www.examsempire.com/>**

**Product Version**

- 1. Up to Date products, reliable and verified.**
- 2. Questions and Answers in PDF Format.**



**<https://examsempire.com/>**

# Latest Version: 6.0

## Question: 1

The patient has \_\_\_\_.



- A. Veneers.
- B. A wisdom tooth that is extremely crooked.
- C. Four wisdom teeth.
- D. Two wisdom teeth.

**Answer: A**

Explanation:

The patient has veneers. Veneers are thin shells, typically made of porcelain or resin, that cover the front surface of the teeth to improve their appearance. These are often used to treat teeth that are discolored, chipped, worn down, uneven, or abnormally spaced. The x-ray of the patient shows that in addition to veneers, the patient has undergone other extensive dental work.

The x-ray indicates the presence of crowns, which are caps placed over teeth to restore their shape, size, strength, and appearance. Bridges, which are used to replace missing teeth by anchoring onto adjacent teeth, are also visible. The presence of dental implants suggests that some teeth were missing and have been replaced with these permanent fixtures that are anchored directly into the jawbone.

Moreover, the x-ray shows missing teeth, which could be due to various reasons such as decay, injury, or even prior extractions for orthodontic reasons. Root canals visible in the x-ray indicate that the patient has had treatment to remove infected or damaged tissue from within the tooth, thereby saving the tooth from extraction.

In terms of wisdom teeth, the description mentions different scenarios: a wisdom tooth that is extremely crooked, four wisdom teeth, and two wisdom teeth. This suggests variability in the number

and condition of wisdom teeth among different patients or possibly different interpretations of a single patient's x-ray over time. Wisdom teeth, also known as third molars, often cause issues such as crowding, infections, or impactions, leading to the need for their extraction.

In summary, the x-ray shows a comprehensive range of dental treatments indicating significant attention to dental health and restoration. The combination of veneers, crowns, bridges, implants, and root canals along with the management of wisdom teeth illustrates a complex dental history aimed at improving oral function and aesthetics.

## Question: 2

A patient could suffer from bleeding complications if he or she has a(n) \_\_\_\_\_ condition.

- A. Lung.
- B. Infectious.
- C. Eye.
- D. Kidney.

**Answer: D**

Explanation:

A patient could suffer from bleeding complications if he or she has a(n) \_\_\_\_\_ condition. \*PLung.\*  
Kidney. Patients are at a higher risk of bleeding complications if they have kidney problems. Kidney diseases can affect the body's ability to clot blood properly, leading to an increased risk of bleeding. This is because the kidneys play a critical role in producing substances that help in the blood clotting process and in metabolizing and clearing certain medications that can impact bleeding risk.

Liver. Similarly, liver conditions can also lead to an increased risk of bleeding complications. The liver produces proteins necessary for blood clotting, and liver dysfunction can result in a deficiency of these clotting factors, thereby increasing the risk of bleeding. Furthermore, liver disease can lead to portal hypertension, which can cause varices (enlarged veins) that may bleed.

Infectious. Infectious conditions, depending on their nature and location, could potentially lead to bleeding issues. For example, infections that involve the gastrointestinal tract, such as those caused by certain bacteria or viruses, can lead to ulcerations and subsequently to bleeding. Additionally, some systemic infections can cause disseminated intravascular coagulation (DIC), a serious condition that leads to widespread clotting and bleeding in the blood vessels.

Eye. While less common in terms of systemic bleeding risks, certain eye conditions can lead to localized bleeding. For example, diabetic retinopathy, a complication of diabetes, can cause abnormal blood vessel growth in the retina, which can bleed. However, eye conditions generally do not contribute to systemic bleeding risks unless they are part of a broader systemic disease process.

Kidney. Reiterating the specific risks associated with kidney disease, it is crucial for patients with kidney dysfunction to be closely monitored for signs of bleeding, especially if they are undergoing surgical procedures or are on medication regimens that could further impair clotting capabilities. Management of bleeding risk in kidney disease often involves careful adjustment of medications, strict control of blood pressure, and regular blood tests to monitor kidney function and clotting status.

## Question: 3

Which of the following is NOT true of the trays used for in-office topical fluoride application?

- A. They are disposable.
- B. They are available in primary, mixed, and adult dentition sizes.
- C. They should be custom-fitted.
- D. They should be used only once.

**Answer: C**

Explanation:

Correct Answer: They should be custom-fitted.

The statement that trays used for in-office topical fluoride application should be custom-fitted is not true. In reality, these trays are not custom-made to fit the individual patient's mouth. Instead, they are pre-manufactured in various standard sizes that are designed to accommodate a range of mouth sizes and shapes. The primary objective is to ensure that the trays can cover the dental arches adequately for effective fluoride application, without the need for each tray to be uniquely tailored to the patient.

These fluoride application trays are disposable, which means they are designed for single use only. This practice helps maintain high hygiene standards by reducing the risk of cross-contamination between patients. Since the trays are not reused, there is no requirement for sterilization between uses, which can be both time-consuming and resource-intensive in a dental practice setting.

The trays are available in different sizes, which allows the dental practitioner to choose the most appropriate size for each patient. This selection process is important to ensure that the fluoride gel or foam applied in the tray contacts all necessary surfaces of the teeth effectively. Proper coverage is crucial for the fluoride treatment to be effective in strengthening the tooth enamel and helping to prevent cavities.

In conclusion, while fluoride application trays are standard in size and disposable, they are not custom-fitted for each patient. The confusion might arise from the use of other dental devices like mouth guards or bleaching trays, which often do require custom fitting. However, for the purpose of fluoride application in a professional setting, pre-sized, disposable trays are the norm.

### Question: 4

Which of the following would you use when taking a dental impression to prevent the flow of impression material into the back of the throat?

- A. Sticky wax.
- B. Utility wax.
- C. Base plate wax.
- D. Bite registration wax.

**Answer: B**

Explanation:

The correct choice for preventing the flow of impression material into the back of the throat during a dental impression is utility wax. Utility wax is notably pliable at room temperature, which makes it highly adaptable for various uses in dental procedures. Its pliability allows it to be easily molded to fit specific

areas and contours within the mouth. This characteristic is particularly useful in creating a barrier or a seal around the periphery of the impression trays.

When taking dental impressions, especially of the upper arch, there is a risk of impression material flowing towards the back of the throat, which can cause discomfort or gagging in patients. To prevent this, utility wax can be applied to the posterior edge of the impression tray. The wax forms an effective barrier that blocks the pathway of the impression material towards the throat. By doing so, it not only enhances patient comfort but also increases the safety of the impression-taking process.

Moreover, utility wax can also protect soft tissues from potential damage. During the impression process, the edges of the impression trays or the hardness of the impression materials can sometimes irritate or harm the soft tissues of the mouth, such as the gums and cheeks. Applying utility wax to the tray edges softens these contact points, significantly reducing the risk of tissue abrasion or discomfort. In comparison to other types of dental waxes like sticky wax, base plate wax, or bite registration wax, utility wax is generally more suited for this specific purpose due to its superior pliability and ease of application. Sticky wax, for instance, is used more for temporarily holding components together in prosthetics and is not as pliable. Base plate wax is harder and used primarily for creating denture bases during the preliminary stages of denture fabrication, making it less ideal for forming a soft, protective barrier. Bite registration wax, although used in the impression process, is designed specifically for recording the occlusal relationships between the upper and lower teeth, rather than for providing a throat barrier.

Therefore, in dental practice, when it comes to choosing a material to prevent the flow of impression material into the back of the throat, utility wax is the most effective and appropriate selection due to its physical properties and ease of use.

### Question: 5

Using a petroleum based product immediately before putting on exam gloves will cause which of the following to gloves?

- A. Breakdown.
- B. Tearing.
- C. Preserve integrity.
- D. Strengthen.

**Answer: A**

Explanation:

Using a petroleum-based product, such as lotions or ointments that contain petroleum jelly before putting on exam gloves, especially those made from latex or vinyl, can have detrimental effects on the material of the gloves. Petroleum products have a degrading effect on these materials, leading to a breakdown of the glove's structural integrity.

The breakdown occurs because the petroleum reacts chemically with the latex or vinyl, causing the material to weaken. This weakening manifests as a loss of elasticity and the formation of micro-tears in the glove material. As the glove material becomes compromised, tiny holes may develop, which are often imperceptible at first. This degradation reduces the glove's effectiveness in providing a barrier against contaminants or pathogens.

As a result of this breakdown, the gloves become more susceptible to tearing or ripping during use. This can occur during routine tasks such as pulling on the gloves or during medical procedures that require

manual dexterity and apply stress to the material. Once torn or ripped, the gloves fail to provide the necessary barrier protection between the healthcare worker's skin and environmental contaminants, including bodily fluids or chemical agents.

It is critical, therefore, to avoid the use of petroleum-based products on the hands immediately before using latex or vinyl gloves. Alternative non-petroleum-based hand moisturizers should be considered, especially those that are labeled as compatible with latex or vinyl gloves. Additionally, ensuring hands are dry before donning gloves can also help preserve the integrity and effectiveness of the gloves in maintaining a safe and hygienic barrier during medical examinations or procedures.

### Question: 6

When do the deciduous teeth start developing?

- A. During infancy.
- B. During early childhood.
- C. Before age twelve.
- D. In the fetus.

**Answer: D**

Explanation:

The correct answer to when deciduous teeth start developing is "In the fetus." Deciduous teeth, also commonly known as baby teeth or primary teeth, begin their development in utero, specifically during the embryonic stage of pregnancy. This initial phase of tooth development is crucial as it lays the foundation for proper oral health and the alignment of permanent teeth later in life.

Tooth development, or odontogenesis, is a complex process that starts as early as six weeks into pregnancy. At this stage, the basic structures of the future teeth are formed through a series of growth and differentiation stages involving the dental lamina and tooth buds. These structures gradually develop into the hard tissues and distinctive shapes of teeth. By the time a baby is born, the crowns of the primary teeth are almost fully formed, although the teeth themselves do not typically begin to emerge through the gums until the infant is between six months to one year old.

Interestingly, not only the primary teeth but also the beginnings of the permanent teeth form during fetal development. The tooth buds for the permanent teeth start to develop behind the primary teeth around the fourth month of gestation. This fact highlights the importance of maternal nutrition and health during pregnancy, as it can significantly influence the development and mineralization of the teeth.

In conclusion, the development of deciduous teeth starts much earlier than many might assume, occurring during the fetal stage. This early initiation is essential to ensure that children have healthy and well-formed teeth, which are vital for proper nutrition through the ability to chew food, as well as for the development of clear speech.

### Question: 7

Which of the following would you choose to clean an amalgam restoration?

- A. Greenie.

- B. Brownie.
- C. Air abrasion unit.
- D. Diamond disk.

**Answer: B**

Explanation:

When considering the best tool to clean an amalgam restoration, it is important to understand the properties of amalgam and the effects of different cleaning tools on this material. Amalgam, used for dental fillings, is a mixture of metals including mercury, silver, tin, and copper which hardens into a durable material. The choice of cleaning tool should effectively clean without damaging the amalgam surface.

The options listed for cleaning amalgam restorations are Greenie, Brownie, Air abrasion unit, and Diamond disk. Each of these tools has distinct characteristics and uses in dental procedures. Starting with the Greenie, it is a mild abrasive tool typically used for polishing composites rather than for cleaning amalgam. It is softer and less abrasive than a Brownie, making it less suitable for amalgam which might need a more robust cleaning to remove surface oxidation or stains.

The Brownie, on the other hand, is a moderately abrasive tool ideal for cleaning and polishing amalgam restorations. It is designed to remove tarnish and restore luster without excessively removing material from the restoration itself. When using a Brownie, it is important to apply light pressure and use a slow speed to avoid overheating the amalgam. The surface should appear to have a dull luster after use, indicating that it has been effectively cleaned without being damaged. It is crucial to change the tips of the Brownie before they are worn down to the metal shank, which can scratch and damage the amalgam surface.

The Air abrasion unit is another option, which uses a fine stream of abrasive particles to clean and prepare teeth surfaces. However, for cleaning amalgam restorations, this method may be too aggressive, potentially removing too much material and damaging the integrity of the filling. Lastly, the Diamond disk is a highly abrasive tool generally used for cutting and shaping hard materials like enamel or old restorations but is too abrasive for cleaning amalgam restorations. Its aggressive nature can lead to excessive removal of the amalgam material, potentially weakening the restoration. Based on these considerations, the Brownie is the most appropriate choice for cleaning an amalgam restoration. It effectively cleans the surface with a controlled level of abrasiveness that minimizes the risk of damage. Proper technique, such as using light pressure and a slow speed, along with timely replacement of the tool before it is overly worn, is crucial to avoid scratching or otherwise damaging the amalgam.

### Question: 8

The visible light used for curing sealants is:

- A. Infrared light.
- B. Blue spectrum.
- C. Red spectrum.
- D. Orange spectrum.

**Answer: B**

Explanation:

The correct answer to the question regarding the type of visible light used for curing sealants is "Blue spectrum."

Blue light is commonly used in various dental and industrial applications for curing photopolymer resins, including sealants. This specific spectrum of light is highly effective in initiating the polymerization (curing) process of the materials, which solidifies them quickly and effectively. The most typical wavelength range for this purpose falls between 400 to 500 nanometers, which is why it is referred to as "blue spectrum."

The reason blue light is preferred over other spectrums, such as red or orange, is due to its optimal energy level that matches the activation energy required by the photoinitiators present in many sealants and resins. Photoinitiators are chemical compounds that react to specific wavelengths of light to begin the curing process. Blue light provides the right balance of energy to activate these compounds efficiently without causing damage or excessive heat to the surrounding material.

It's also worth noting that while most modern curing lights use blue spectrum visible light, some use ultraviolet (UV) light. UV light units, operating typically at wavelengths below 400 nanometers, are also effective in initiating the polymerization of certain materials. However, due to concerns about UV exposure and its potential health risks, blue light units are more commonly used, particularly in settings requiring frequent or prolonged use, such as in dental practices.

In summary, blue spectrum light is the standard and most common choice for curing sealants due to its effectiveness in activating photoinitiators and its safer profile compared to UV light. This makes it an essential tool in both dental and various

### Question: 9

When should contaminated items be soaked?

- A. After sterilization.
- B. Immediately.
- C. Before washing.
- D. Before drying.

**Answer: B**

Explanation:

Contaminated items should be soaked immediately after use. This is a critical step in infection control protocols within medical, laboratory, and similar settings where equipment and instruments come into contact with biological materials. Immediate soaking minimizes the risk of biofilm formation and the hardening of organic materials, such as blood or tissue, which can be more difficult to remove if allowed to dry.

Soaking the items right away in an appropriate decontaminating solution helps to loosen and dissolve these substances, making the subsequent cleaning processes more effective. The type of solution used for soaking might vary depending on the nature of the contamination and the material of the contaminated items. Commonly used solutions include enzymatic cleaners, which are particularly effective at breaking down organic matter.

The importance of immediate soaking extends beyond just cleaning efficacy. It also plays a significant role in the safety of the staff handling the items. By preventing the drying and hardening of potentially



infectious material, immediate soaking reduces the risk of aerosolization of harmful particles during later cleaning stages. This can significantly decrease the likelihood of occupational exposure to infectious agents.

Furthermore, immediate soaking is essential for the maintenance of the instruments themselves. Materials that remain on instruments for an extended period can cause corrosion or other damage, potentially shortening the lifespan of expensive equipment. Proper immediate care and maintenance can result in significant cost savings over time.

In summary, contaminated items must be soaked immediately to ensure effective cleaning, maintain safety standards, protect staff health, and prolong the usability of instruments. This step is essential before washing, drying, and especially before any sterilization processes take place.

## Question: 10

Topical anesthetics applied to the skin can cause what kind of inflammatory condition?

- A. Contact dermatitis.
- B. Abrasions.
- C. Bruising.
- D. Edema.

**Answer: A**

Explanation:

Topical anesthetics are medications applied to the skin to temporarily numb and relieve pain in a specific area. While they are effective for their intended purpose, they can sometimes lead to an inflammatory skin condition known as contact dermatitis.

Contact dermatitis is a form of inflammation of the skin that arises from direct contact with irritants or allergens. In the case of topical anesthetics, the active ingredients or even preservatives and other chemical components in the formulation can act as potential irritants or allergens. Symptoms of contact dermatitis include redness, itching, swelling, and in severe cases, blistering and peeling of the skin.

The mechanism behind contact dermatitis involves an immune response triggered by the skin's contact with the offending substance. This response can be categorized into two types: irritant contact dermatitis and allergic contact dermatitis. Irritant contact dermatitis is more common and occurs as a direct result of the anesthetic damaging the skin barrier. Allergic contact dermatitis, on the other hand, is an immune-mediated response where the body's immune system reacts to a substance as though it were harmful, even though it might not be.

It is important to differentiate contact dermatitis from other skin conditions that might arise in different contexts. Abrasions and bruising, which are physical damages to the skin, are typically caused by trauma and do not stem from a chemical reaction or immune response. Edema, characterized by the accumulation of fluid in the tissues leading to swelling, commonly results from circulatory problems, rather than direct skin contact with chemicals.

In summary, when using topical anesthetics, it is crucial to be aware of the potential for developing contact dermatitis. Users should monitor their skin's reaction to these products and seek medical advice if signs of inflammation occur. Preventive measures include testing the product on a small skin area before widespread application and consulting healthcare providers about any known allergies or sensitivities to ingredients in topical anesthetics.

**Thank You for Trying Our Product**

**Special 16 USD Discount Coupon: NSZUBG3X**

**Email:** [support@examsempire.com](mailto:support@examsempire.com)

**Check our Customer Testimonials and ratings  
available on every product page.**

**Visit our website.**

**<https://examsempire.com/>**